



April 5, 2010

Submission Request c/o Joan McClure  
National Comprehensive Cancer Network  
275 Commerce Dr, Suite 300  
Fort Washington, PA 19034

**RE: Request for Addition of Oforta (oral fludarabine phosphate) for Use in 1<sup>st</sup> Line CLL and 1<sup>st</sup> and 2<sup>nd</sup> Line NHL**

Name: Julia Petses  
Company/Organization: sanofi-aventis US  
Address: 55 Corporate Dr. Bridgewater, NJ 08807  
Phone: (908) 981-7287  
Email: julia.petses@sanofi-aventis.com  
Date of request: April 5, 2010  
NCCN Guidelines Panel: NHL

Dear Ms. McClure:

As the National Comprehensive Cancer Network (NCCN) non-Hodgkin's lymphoma (NHL) Panel reviews the NCCN Clinical Practice Guidelines in Oncology for NHL, we respectfully request consideration of the inclusion of Oforta™ (oral fludarabine phosphate) as a treatment option for first-line chronic lymphocytic leukemia (CLL) and first- and second-line NHL.

**Request for Addition of Oforta (oral fludarabine phosphate) for Use in 1<sup>st</sup> Line CLL and 1<sup>st</sup> and 2<sup>nd</sup> Line NHL**

Oforta (oral fludarabine phosphate) is approved for use in adult patients with B-cell CLL whose disease has not responded to or has progressed during or after treatment with at least one standard alkylating-agent containing regimen.<sup>1</sup> Additionally, data exists evaluating the use of oral fludarabine phosphate as a single agent in the treatment of first line (previously untreated) CLL, including two studies summarized within the Oforta product labeling.<sup>2-5</sup> Patients received oral fludarabine phosphate 40 mg/m<sup>2</sup> daily for 5 days, every 28 days in both studies.<sup>2-4</sup> In Study 2 of the Oforta product labeling, the overall response rate according to NCI criteria was 80.2%, and 71.6% according to IWCLL criteria.<sup>2</sup> Myelosuppression was the most frequent grade 3/4 toxicity (granulocytopenia, 32.1%; anemia, 9.9%; thrombocytopenia, 4.9%). In Study 3, the overall response rate according to modified NCI criteria was 74%.<sup>3-4</sup> There was no significant difference in main toxicities between oral and IV fludarabine in this study. Data also exists for the use of oral fludarabine phosphate (24-30 mg/m<sup>2</sup>) plus oral cyclophosphamide (150-250 mg/m<sup>2</sup>) given for 3-5 days every 4 weeks in previously untreated CLL patients that demonstrate efficacy and acceptable toxicity.<sup>3,6-8</sup> The use of oral fludarabine phosphate in patients with NHL has also been evaluated, and includes previously treated and untreated patients with small lymphocytic, lymphoplasmacytic, follicular, marginal zone (including MALT lymphoma), and mantle cell lymphoma.<sup>9-12</sup> These data include the use of oral fludarabine phosphate (40 mg/m<sup>2</sup>) as a single agent<sup>9</sup>, oral fludarabine phosphate (25 mg/m<sup>2</sup>) plus oral cyclophosphamide (150 mg/m<sup>2</sup>)<sup>10</sup>, as well as oral fludarabine phosphate (40 mg/m<sup>2</sup>) plus intravenous rituximab.<sup>11-12</sup>

**Specific change recommended**

Consider the addition of Oforta (or listed Oforta-containing regimen) as a suggested treatment regimen in the following:

- CLL: First-line single agent therapy<sup>2-5</sup> and combined with oral cyclophosphamide<sup>6-8</sup> (CSLL-D)
- Follicular Lymphoma: First-line therapy combined with oral cyclophosphamide<sup>10</sup> (low dose regimen in elderly patients) or combined with rituximab<sup>11</sup> (FOLL-B)
- Follicular Lymphoma: Second-line and subsequent therapy as single agent<sup>9</sup> and in combination with rituximab<sup>12</sup> (FOLL-B)
- Mantle Cell Lymphoma: Second-line single agent therapy<sup>9</sup> (MANT-A)

## FDA Status

Oforta (oral fludarabine phosphate) is approved for use in adult patients with B-cell CLL whose disease has not responded to or has progressed during or after treatment with at least one standard alkylating-agent containing regimen,<sup>1</sup> and is not approved for use in 1<sup>st</sup> line CLL or 1<sup>st</sup> or 2<sup>nd</sup> line NHL.

## Rationale for recommended change

Multiple phase II studies evaluating the use of oral fludarabine phosphate as a single agent and in combination with oral cyclophosphamide or rituximab provide evidence for use in patients with CLL and NHL by demonstrating efficacy and acceptable toxicity.

## Literature support

1. Oforta [package insert]. Bridgewater, NJ: sanofi-aventis U.S.; 2009.
2. Rossi J-F, Van Hoof A, De Boeck K, et al. Efficacy and safety of oral fludarabine phosphate in previously untreated patients with chronic lymphocytic leukemia. *J Clin Oncol*. 2004; 22(7):1260-1267.
3. Catovsky D, Richards S, Matutes E, et al. Assessment of fludarabine plus cyclophosphamide for patients with chronic lymphocytic leukaemia (the LRF CLL4 trial): a randomised controlled trial. *Lancet*. 2007; 370:230-239.
4. Hillmen P, Richards S, Catovsky D. Comparison of oral and intravenous fludarabine in the LRF CLL4 trial. *Blood*. 2005;2006: Abstract 722.
5. Friedberg JW, Dong DA, Li S, et al. Oral fludarabine has significant activity in patients with previously untreated chronic lymphocytic leukemia, and leads to increased STAT1 levels in vivo. *Leukemia Res*. 2004;28:139-147.
6. Cazin B, Divine M, Lepretre S, et al. High efficacy with five days schedule of oral fludarabine phosphate and cyclophosphamide in patients with previously untreated chronic lymphocytic leukaemia. *Br J Haematol*. 2008;143, 54-59.
7. Fabbri A, Lenoci M, Gozzetti A, et al. Low-dose fludarabine plus cyclophosphamide in elderly patients with chronic lymphoproliferative disorder. *Hematol J*. 2004;5:472-474.
8. Laurenti L, Tarnani M, De Padua L, et al. Oral fludarabine and cyclophosphamide as front-line chemotherapy in patients with chronic lymphocytic leukemia. The impact of biological parameters in the response duration. *Ann Hematol*. 2008;87:891-898.
9. Ogawa Y, Hotta T, Tobinai, et al. Phase I and pharmacokinetic study of oral fludarabine phosphate in relapse indolent B-cell non-Hodgkins' lymphoma. *Ann Oncol*. 2006;17:330-333.
10. Fabbri A, Lenoci M, Gozzetti A, et al. Low-dose oral fludarabine plus cyclophosphamide as first-line treatment in elderly patients with indolent non-Hodgkin lymphoma. *Br J Haematol*. 2007;139:90-93.
11. Salar A, Domingo-Domenech E, Estany C, et al. Combination therapy with rituximab and intravenous or oral fludarabine in the first-line, systemic treatment of patients with extranodal marginal zone b-cell lymphoma of the mucosa-associated lymphoid tissue type. *Cancer*. 2009;1-8.
12. Tobinai K, Ishizawa K, Ogura M, et al. Phase II study of oral fludarabine in combination with rituximab for relapsed indolent B-cell non-Hodgkin lymphoma. *Cancer Sci*. 2009;100:1951-1956.