

Guideline Page and Request	Panel Discussion/References	Institution Vote			
		YES	NO	ABSTAIN	ABSENT
<p>MGF-1 Internal request: In response to the FDA approval of trilaciclib as an option to reduce the frequency of chemotherapy-induced bone marrow suppression in adults receiving certain types of chemotherapy for extensive-stage small cell lung cancer. The panel requested the addition of a footnote on the MGF-1 stating, "Trilaciclib may be used as a prophylactic option to decrease the incidence of chemotherapy-induced myelosuppression when administered before (or G-CSF may be administered after) platinum/etoposide ± immune checkpoint inhibitor-containing regimens or a topotecan-containing regimen for extensive-stage small cell lung cancer (ES-SCLC)."</p> <p>MGF-1 External Submission Submission from G1 Therapeutics (02/12/21) requesting the addition of trilaciclib as a footnote on MGF-1 (Prior to First Chemotherapy Cycle) as a preferred option to decrease the incidence of chemotherapy induced myelosuppression from regimens containing platinum + etoposide +/- checkpoint inhibition; or a regimen containing topotecan.</p>	<p>Based on the review of the data in the noted references and the recent FDA approval, the panel consensus was to include trilaciclib as a footnote on MGF-1. This is a category 2B recommendation.</p> <ul style="list-style-type: none"> • See submission for references 	12	3	3	14
<p>MGF-3 Internal request: In response to the FDA approval of trilaciclib as an option to reduce the frequency of chemotherapy-induced bone marrow suppression in adults receiving certain types of chemotherapy for extensive-stage small cell lung cancer. The panel requested the addition of a footnote on the MGF-3 stating, "Trilaciclib may be used as a prophylactic option to decrease the incidence of chemotherapy-induced myelosuppression when administered before (or G-CSF may be administered after) platinum/etoposide ± immune checkpoint inhibitor-containing regimens or a topotecan-containing regimen for extensive-stage small cell lung cancer (ES-SCLC)."</p>	<p>Based on the review of the data in the noted references and the recent FDA approval, the panel consensus was to include trilaciclib as a footnote on MGF-3. This is a category 2B recommendation.</p> <ul style="list-style-type: none"> • See submission for references 	11	4	3	14

<p>MGF-3 External Submission Submission from G1 Therapeutics (02/12/21) requesting the addition of trilaciclib as a footnote on MGF-3 (Prior to Second and Subsequent Chemotherapy Cycles) as a preferred option to decrease the incidence of chemotherapy induced myelosuppression from regimens containing platinum + etoposide +/- checkpoint inhibition; or a regimen containing topotecan.</p>					
<p>MGF-A 1 of 5 and 2 of 5 Internal request: In response to the FDA approval of trilaciclib as an option to reduce the frequency of chemotherapy-induced bone marrow suppression in adults receiving certain types of chemotherapy for extensive-stage small cell lung cancer. The panel requested the addition of a footnote on the MGF-A, 1 and 2 of 5, stating, “Trilaciclib may be used as a prophylactic option to decrease the incidence of chemotherapy-induced myelosuppression when administered before (or G-CSF may be administered after) platinum/etoposide ± immune checkpoint inhibitor-containing regimens or a topotecan-containing regimen for extensive-stage small cell lung cancer (ES-SCLC).”</p> <p>MGF-A (1 of 5 and 2 of 5) External Submission Submission from G1 Therapeutics (02/12/21) requesting the addition of trilaciclib as footnotes on MGFA, 1&2/5 (Under Small Cell Lung Cancer regimens) as a preferred option to decrease the incidence of chemotherapy induced myelosuppression from regimens containing platinum + etoposide +/- checkpoint inhibition; or a regimen containing topotecan.</p>	<p>Based on the review of the data in the noted references and the recent FDA approval, the panel consensus was to include trilaciclib as a footnote on MGF-A, 1 of 5 and 2 of 5.</p> <ul style="list-style-type: none"> • See submission for references 	14	1	3	14
<p>ANEM-1 Internal request: In response to the FDA approval of trilaciclib as an option to reduce the frequency of chemotherapy-induced bone marrow suppression in adults receiving certain types of chemotherapy for extensive-stage small cell lung cancer. The panel requested the addition of a footnote on the ANEM-1 stating, “Trilaciclib may be used as a prophylactic option to</p>	<p>Based on the review of the data in the noted references and the recent FDA approval, the panel consensus was to include trilaciclib as a footnote on ANEM-1. Based on the review of the data in the noted references and the recent FDA approval, the panel consensus was to include trilaciclib as a footnote on MGF-1. This is a category 2B recommendation.</p>	9	6	3	14

<p>decrease the incidence of chemotherapy-induced myelosuppression when administered before (or G-CSF may be administered after) platinum/etoposide ± immune checkpoint inhibitor-containing regimens or a topotecan-containing regimen for extensive-stage small cell lung cancer (ES-SCLC).”</p> <p>ANEM-1 External Submission Submission from G1 Therapeutics (02/12/21) requesting the addition of trilaciclib as a new pathway in ANEM-1 as preferred option to decrease the incidence of chemotherapy induced myelosuppression from regimens containing platinum + etoposide +/- checkpoint inhibition; or a regimen containing topotecan.</p>	<ul style="list-style-type: none"> • See submission for references 				
--	---	--	--	--	--