## NCCN Guidelines for Bladder Cancer V.2.2019 – Web Teleconference on 03/01/19

Guideline Page and Request	Panel Discussion/References	Institution Vote			
and Request		YES	NO	ABSTAIN	ABSENT
BL-F Internal request:					
Panel discussion to consider in the context of a BCG shortage the inclusion of intravesical epirubicin for induction therapy.	The panel consensus was to include intravesical epirubicin as an option for induction therapy when there is a BCG shortage. This is a category 2A recommendation.	15	0	0	17
BL-F Internal request:					
Panel discussion to consider in the context of a BCG shortage the inclusion of intravesical valrubicin for induction therapy.	The panel consensus was to include intravesical valrubicin as an option for induction therapy when there is a BCG shortage. This is a category 2A recommendation.	15	0	0	17
BL-F Internal request:					
Panel discussion to consider in the context of a BCG shortage the inclusion of intravesical docetaxel for induction therapy.	The panel consensus was to include intravesical docetaxel as an option for induction therapy when there is a BCG shortage. This is a category 2A recommendation.	15	0	0	17
BL-F Internal request:					
Panel discussion to consider in the context of a BCG shortage the inclusion of intravesical gemcitabine/docetaxel for induction therapy.	The panel consensus was to include intravesical gemcitabine/docetaxel as an option for induction therapy when there is a BCG shortage. This is a category 2A recommendation.	15	0	0	17
BL-F Internal request:					
Panel discussion to consider in the context of a BCG shortage the inclusion of intravesical gemcitabine/mitomycin for induction therapy.	The panel consensus was to include intravesical gemcitabine/mitomycin as an option for induction therapy when there is a BCG shortage. This is a category 2A recommendation.	15	0	0	17