Name: Suzana Giffin, AVP

Company/Organization: Merck & Co., Inc.

Address: 2000 Galloping Hill Rd. Kenilworth, NJ 07033

Phone: 908-740-6708

Email: suzana.giffin@merck.com Date of Request: April 29, 2020

NCCN Guidelines Panel: Head and Neck Cancers Panel

NCCN Head and Neck Cancers Panel: On behalf of Merck & Co., Inc., I respectfully request that the NCCN Head and Neck Cancers Panel review the enclosed information for KEYTRUDA® (pembrolizumab), in reference to updated dosing recommendations for pembrolizumab.

<u>Specific Changes</u>: We respectfully request the inclusion of the updated dosing recommendations for pembrolizumab, either 200 mg every 3 weeks or 400 mg every 6 weeks administered as a 30-minute intravenous (IV) infusion until disease progression, unacceptable toxicity, or up to 24 months for the treatment of adult patients with head and neck squamous cell carcinoma (HNSCC), to SYST-A (page 2 of 6), SYST-A (page 3 of 6), and pages MS-37, MS-38, MS-40, and MS-41 in the NCCN Head and Neck Cancers Guidelines.

FDA Clearance:

Adult Indications: Additional Dosing Regimen of 400 mg Every 6 Weeks

 KEYTRUDA is indicated for use at an additional recommended dosage of 400 mg every 6 weeks for all approved adult indications. This indication is approved under accelerated approval based on pharmacokinetic data, the relationship of exposure to efficacy, and the relationship of exposure to safety. Continued approval for this dosing may be contingent upon verification and description of clinical benefit in the confirmatory trials.

Head and Neck Squamous Cell Cancer

- KEYTRUDA, in combination with platinum and fluorouracil (FU), is indicated for the first-line treatment of patients with metastatic or with unresectable, recurrent HNSCC.
- KEYTRUDA, as a single agent, is indicated for the first-line treatment of patients with metastatic or with unresectable, recurrent HNSCC whose tumors express PD-L1 [Combined Positive Score (CPS) ≥1] as determined by an FDA-approved test.
- KEYTRUDA, as a single agent, is indicated for the treatment of patients with recurrent or metastatic HNSCC with disease progression on or after platinum-containing chemotherapy.

Please refer to the KEYTRUDA (pembrolizumab) prescribing information for other FDA-approved indications.¹

<u>Rationale</u>: KEYTRUDA (pembrolizumab) is now FDA-approved for use at an additional recommended dosage of 400 mg every 6 weeks for all approved adult indications. This indication is approved under accelerated approval based on pharmacokinetic data, the relationship of exposure to efficacy, and the relationship of exposure to safety. Continued approval for this dosing may be contingent upon verification and description of clinical benefit in the confirmatory trials. Based on the modeling of

dose/exposure efficacy and safety relationships and observed pharmacokinetic data from an interim analysis of 41 patients with melanoma treated with pembrolizumab 400 mg every 6 weeks, there are no anticipated clinically significant differences in efficacy and safety between pembrolizumab doses of 200 mg or 2 mg/kg every 3 weeks or 400 mg every 6 weeks.¹ Additionally, this submission includes the pharmacokinetic modeling and simulation data published by Lala et al.² and the preliminary pharmacokinetic, efficacy and safety results from the KEYNOTE-555 trial (investigating a Q6W [every 6 weeks] dosing regimen for pembrolizumab in patients with advanced melanoma [NCT03665597]),³ which support the recent label change.

The following resources are submitted to assist the committee with their review.

- 1. KEYTRUDA (pembrolizumab) prescribing information. Merck & Co., Inc.
- 2. Lala M, Li TR, De Alwis DP, et al. A six-weekly dosing schedule for pembrolizumab in patients with cancer based on evaluation using modelling and simulation. *European Journal of Cancer*. 2020;131:68-75. doi: 10.1016/j.ejca.2020.02.016 [Epub ahead of print].
- Lala M, Akala O, Chartash E, et al. Pembrolizumab 400 mg Q6W dosing: First clinical outcomes data from KEYNOTE-555 cohort B in patients with metastatic melanoma. Presented at the American Association for Cancer Research (AACR) Annual Meeting; April 27-28, 2020; Virtual Meeting.

Thank you for considering this request. Below is my contact information should you need to contact me for additional information.

Sincerely,

Suzana Giffin, AVP

Merck & Co., Inc. 2000 Galloping Hill Rd Kenilworth, NJ 07033

908-740-6708

suzana.giffin@merck.com